

Symposium on Advanced Wound Care (SAWC) Spring 2024 | Orlando, FL

BACKGROUND

Nursing shortages have system-wide domino effects, 1,2 starting in the acute hospital settings and continuing on to home health care. Professional nurses' positive impact on patient safety, including outcomes in wound care, is well documented. The lack of sufficient nurses working in home care has at times delayed hospitalized patients from being discharged, and/or posed significant time constraints to home care nurses who need to perform frequent wound care utilizing conventional wound care products. Today's healthcare challenges demand new approaches beyond the prevailing models to increase nursing productivity and still ensure excellent patient care and outcomes.

METHODS

A sample population of five established home care patients with challenging wounds, requiring two or more nursing visits a week were treated with transforming powder dressing (TPD*) for evaluation. Our objective was to decrease the time and frequency of home nursing visits for wound care without compromising patient outcomes or increasing costs. A home health nurse reviewed multiple established home care patients who had challenging wounds from a variety of etiologies and were being visited more than once weekly, and who were not significantly improving with standard of care (SOC).

Transforming powder dressing (TPD) is a commercially available dressing comprised primarily of polymers similar to those used in contact lenses. When hydrated with saline, TPD granules aggregate to form a moist, oxygen-permeable barrier that covers and protects the wound while releasing excess exudate through vapor transpiration. It may be left in on the wound for up to 30 days and topped off as needed without requiring primary dressing changes.

DISCUSSION

Strategic partnering with industry to trial innovations and evolve best practices resulted in improved home-health nursing efficiency. By incorporating TPD, clinical outcomes and patient satisfaction were improved while helping our nursing colleagues become as efficient as possible in this highly challenging healthcare environment.

REFERENCES

- 1. Collins, Rhonda. Reimagining Nursing for the Future: Nurses are Innovators. CNO.3404.Report.202206.INRL_.pdf. Accessed 02JAN2023.
- 2. Jacobs, A. (2022) Nursing is in crisis: Staff shortages put patients at risk. New York Times. 'Nursing Is in Crisis': Staff Shortages Put Patients at Risk-The New York Times (nytimes.com).
- 3. Ross, Jacqueline. Nursing Shortage Creating Patient Safety Concerns. Vol. 37, issue 4, P565-567, Aug. 01, 2022. Published June 30, 2022. DOI: https://doi.org/10.1016/j.jopan.2022.05.078.
- 4. Hall, John. Workforce issues loom over wound care. Nov. 11, 2022. Mcknights.com.

ADVANCING NURSING EFFICIENCY USING A NOVEL EXTENDED-WEAR WOUND DRESSING IN HOME HEALTH CARE AND BEYOND

Jodi Thompson, RN, BSN, CWOCN; Cynthia Hayes, RN, MHA, BSN, Director; Baptist Home Care | Paul Crockett, MHA, BSN, RN: Vice President

RESULTS				Pain	(VAS)	Visits P	er Week
Wound Type	Wound Duration Pre-TPD	Healing Time Post-TPD	Complications Post-TPD	Pre- TPD	Post- TPD	Pre- TPD	Post- TPD
Spider Bite	30 days	42 days	None	10	1	3	1
Radiation Burn	90 days	33 days	None	-	-	7	1
Skin Tear	2 days	44 days	None	10	2	7	1
Amputated Toe	31 days	120 days	None	N/A (neuropathic)		3	1
Lower Leg Ulcer	7 days	Unknown*	None	3	0	3	1
Average	32 days	59.8 days	None	7.7	1	4.6	1

- 82 % reduction in wound assessments / nursing visits per week
- 89% pain reduction for sensate patients post-TPD. Discontinued pain medications.
- No safety issues/complications/adverse effects
- *Wound improved markedly but not quite completely healed when discharged from HH

HOME CARE VISITS / WEEK PAIN SCORES (VAS) Before TPD 4.6 After TPD 1 1 0 PAIN SCORES (VAS)

BROWN RECLUSE SPIDER BITE	CHRONIC RADIATION BURN	TYPE 2 SKIN TEAR	AMPUTATED TOE	LOWER LEG ULCER
PRIOR TO 1 ST TPD APPLICATION 6.5 x 6.0 x 1.0 cm with 2.5 cm tunnel	DAY 0 – PRIOR TO TPD 2 x 1.5 x 0.2 cm	DAY 0 – PRIOR TO TPD 5 x 9 x 0.1 cm	DAY 0 – PRIOR TO TPD 2.2 x 2.5 x 9.5 cm	DAY 0 – PRIOR TO TPD 4.7 x 4.2 x 0.3 cm
DAY 8 – 3 RD VISIT	DAY 15 – 5 TH VISIT	DAY 6 – 2 ND VISIT	DAY 33 – 3 RD VISIT – INITIAL TPD APP	
DATO STORY		DAT 6 - 20 VISIT	Patient Name Date 3 17 23 17 6 20 4 20 Length 7 Width 1.5 Depth 6	DAY 34 – 3 RD VISIT
5.5 x 6.0 x 0.4 cm	1.0 x 1.5 x 0.1 cm	8.5 x 9 x 0.1 cm	0.7 x 1.5 x 0.6	4.7 x 4.2 x 0.3 cm
DAY 22 – 7 TH VISIT	DAY 33 – 9 TH VISIT	DAY 22 – 4 TH VISIT	DAY 42 – 5 TH VISIT	DAY 69 – 4 TH VISIT
		Second of the property of the	Date 4/28/23 1.2 Right Great toe Depth 0.4 Whith D.2 Depth 0.4	ent ID Date 1/10/22
4.0 x 4.5 x 0.3 cm	EPITHELIALIZED	6 x 4 x 0.1 cm	1.2 x 0.2 x 0.4 cm	3.5 x 3.5 x 0.2 cm
DAY 35 – 9 TH VISIT	DAY 42 – 11 TH VISIT	DAY 44	DAY 81	DAY 105
		ATTOCK OF A PART OF STEEL AND A STEEL AND		
3.0 x 3.5 x 0.1 cm	REMAINED HEALED	WOUND HEALED	WOUND HEALED	2 x 2 cm (NO DEPTH)

^{*}Altrazeal Transforming Powder Dressing