
Association of Substance Use and Perinatal Mood and Anxiety **Disorders in Pregnancy**

Introduction


- Perinatal mood and anxiety disorders (PMAD) & substance use disorders (SUD) are associated w/ adverse outcomes in pregnancy^{1,2}
- Suicide and overdose = leading causes of maternal mortality³
- Medicaid population at particularly high risk of both PMAD and SUD⁴
- We aim to assess the relationship of PMAD and SUD in a population of Medicaid-covered births



Literature cited


- . Chang G. Maternal Substance Use: Consequences, Identification, and Interventions. Alcohol Res. 2020 Jun 30;40(2):06.
- 2. Van Niel MS, Payne JL. Perinatal depression: A review. Cleve Clin J Med. 2020 May;87(5):273-277.
- 6;328(21):2159-2161.
- Innovative Screening Approach. Curr Psychiatry Rep. 2020;22(1):1

Erin C. Nacev MD MPH, Kristin C. Prewitt MD MPH, Ann Martinez Acevedo MPH, Maria I. Rodriguez MD MPH Oregon Health & Science University Department of Obstetrics & Gynecology



3. Bruzelius E, Martins SS. US Trends in Drug Overdose Mortality Among Pregnant and Postpartum Persons, 2017-2020. JAMA. 2022 Dec

4. Klawetter S, McNitt C, Hoffman JA, Glaze K, Sward A, Frankel K. Perinatal Depression in Low-Income Women: A Literature Review and



Further information

Please feel free to contact Erin at <u>nacev@ohsu.edu</u> with any thoughts or questions

