

Application of Diamine Silver Fluoride in a 2-Year-Old Patient

Ramirez Correa A, Yáñez Acosta MF, Hernández Rivas MI, Quiroz Leyva CD
University of Guadalajara, Los Altos University Center

Introduction

Introduction: Dental caries disease remains a serious problem worldwide that affects people of all ages, and preventive methods will be the best way to deal with carious lesions (1), and cariostatics such as silver diamine fluoride may be a viable alternative to control the lesions caused by dental caries (2).

Case Report

A 2-year-old male patient came to the pediatric dentistry specialty clinic with the reason that "his front teeth are stained" due to his age and his behavior is a px designated Frankl 4, so it was decided to use FDP at 38%. to stop the progression of caries lesions and to be able to model its behavior as the patient grows and some restorative treatment can be performed. The patient's guardian is instructed on the brushing technique and improving the diet in terms of reducing sugar consumption so that the application of the 38% PDF is successful. In each appointment that the PDF has been applied, 38% have stopped the progression of caries lesions. Which makes it a favorable prognosis for the patient.



D- Second application of FDP at 38% per month.



E- Third application of FDP at 38% per month.

Conclusion

Despite the aesthetic drawback that this material has due to the pigmentation it generates on the tooth surface, FDP is an excellent alternative that we can use in these types of patients to stop the progression of caries lesions, with very good results.



A- Px arrives for consultation.



B- FDP 38% application.



C- Results after application.

BIBLIOGRAFIA

1. Machiulskiene V, Campus G, Carvalho JC, Dige I, Ekstrand KR, Jablonski-Momeni A, et al. Terminology of Dental Caries and Dental Caries Management: Consensus Report of a Workshop Organized by ORCA and Cariology Research Group of IADR. Caries research. 2020;54(1):7-14.
2. Vargas JP, Uribe M, Ortuño D, Verdugo-Paiva F. Silver diamine fluoride compared to atraumatic restorative technique for the treatment of caries in primary and mixed first phase dentition. Medwave. 2020;20(7):e8003.